

# Future changes in Major Stratospheric Warmings in CCMI models

**B. Ayarzagüena<sup>1</sup>, U. Langematz<sup>1</sup>, , L. M. Polvani<sup>2</sup>, J. Abalichin<sup>1</sup>, H. Akiyoshi<sup>3</sup>, A.Klekociuk<sup>4</sup>, M. Michou<sup>5</sup>, O. Morgenstern<sup>6</sup> & L. Oman<sup>7</sup>**


<sup>1</sup>Institut für Meteorologie, Freie Universität Berlin (Germany); <sup>2</sup>Columbia University (USA); <sup>3</sup>NIES, Tsukuba (Japan); <sup>4</sup>Australian Antarctic Division (Australia); <sup>5</sup>Meteo-France, (France); <sup>6</sup>NIWA, (New Zealand); <sup>7</sup>NASA/GSFC (USA)

# Motivation

## Major Stratospheric Warmings (MSWs) in the future

- No clear consensus among recent studies about a possible impact of climate change on MSWs by analyzing chemistry climate models (CCM) simulations [*Butchart et al. 2010; Mitchell et al. 2012*].

Change in SSW freq.  
(1960 -2000 vs 2040-  
2080) in REF-B2 run



[Mitchell et al., 2012]

Possible reason for the uncertainty: competition of different forcings and sensitivity of models to these forcings.

# Goal of the study

**What are the results for the new CCMI runs?**

# Data & models

**REF-C2:** transient run from 1960-2100 under RCP6.0 scenario (GHGs) and A1 scenario (ODS, WMO 2010)

CCMI models	Model resolution	QBO	Solar variability	SSTs
GEOS-CCM	2.5° x 2°, L72 (top:0.01hPa)	Internally generated	No	Prescribed (CESM1)
CNRM-CCM	T42L60 (top: 0.07 hPa)	Internally generated	Yes	Prescribed (CNRM)
NIWA-UKCA	3.75° x 2.5°, L60 (top: 84 km)	Internally generated	No	Coupled to ocean model
CCSRNIES-MIROC 3.2	T42L34 (top: 0.012 hPa)	Nudged	Yes	Prescribed (MIROC 3.2)
EMAC-O	T42L39 (top: 0.01hPa)	Nudged	Yes	Coupled to ocean model
ACCESS	3.75° x 2.5°, L60 (top: 84 km)	Internally generated	No	Prescribed (HadGEM-ES2)

# Mean frequency of MSWs (I)

Periods of study: **PAST** (first 40 winters) and **FUTURE** (last 40 winters)

Model	Past (MSWs/dec)	Future (MSWs/dec)
CCSRNIES-MIROC3.2	2.3	3.0
NIWA-UKCA	0.8 (!)	3.3
GEOS-CCM	1.8	1.8
CNRM-CCM	7.5	7.5
EMAC-O	9.3	7.5
ACCESS	10.0	9.3

## Criterion for MSW identification

- Simultaneous reversal of  $\bar{u}$  at 60°N and 10 hPa &  $\Delta T$  (90°N-60°N) at 10 hPa
- 10 consecutive days of westerly winds after the central date


NCEP/NCAR rean.  
(1960-2000)

# Mean frequency of MSWs (I)

Periods of study: **PAST** (first 40 winters) and **FUTURE** (last 40 winters)

Model	Past (MSWs/dec)	Future (MSWs/dec)
CCSRNIES-MIROC3.2	2.3	3.0
NIWA-UKCA	0.8 (!)	2.2
GEOS-CCM	1.8	1.8
CNRM-CCM	7.5	7.5
EMAC-O	9.3	7.5
ACCESS	10.0	9.3
NCEP/NCAR rean.	5.5	

$\bar{u}$  @ 10 hPa  
NIWA-UKCA



# Mean frequency of MSWs (I)

Periods of study: **PAST** (first 40 winters) and **FUTURE** (last 40 winters)

Model	Past (MSWs/dec)	Future (MSWs/dec)
CCSRNIES-MIROC3.2	2.3	3.0
NIWA-UKCA	0.8 (!)	3.3
GEOS-CCM	1.8	1.8
CNRM-CCM	7.5	7.5
EMAC-O	9.3	7.5
ACCESS	10.0	9.3
NCEP/NCAR rean.	5.5	

## Criterion for MSW identification

- Simultaneous reversal of  $\bar{u}$  at 60°N and 10 hPa &  $\Delta T$  (90°N-60°N) at 10 hPa
- 10 consecutive days of westerly winds after the central date

**No** statistically significant future changes in the mean frequency of MSWs are found except for NIWA-UKCA

## Mean frequency of MSWs (II): Sensitivity of results to different SSW diagnostics

**U6090:** Reversal of  $\bar{u}$  averaged from 60°-90°N  
(Butler et al., under review)

Model	Past (MSWs/dec)	Future (MSWs/dec)
CCSRNIES- MIROC3.2	4.0	5.3
NIWA-UKCA	2.8	3.5
GEOS-CCM	5.5	5.5
CNRM-CCM	10.3	11.8
EMAC-O	8.0	7.3
ACCESS	12.5	11.5
NCEP/NCAR rean.	7.0	

**ZPOL:** Polar-cap averaged 10-hPa Z anomalies  
exceed +3 std (Butler et al. under review)

Model	Past (MSWs/dec)	Future (MSWs/dec)
CCSRNIES- MIROC3.2	3.5	4.0
NIWA-UKCA	4.3	4.5
GEOS-CCM	4.3	5.0
CNRM-CCM	4.0	4.0
EMAC-O	4.8	4.3
ACCESS	3.8	3.8
NCEP/NCAR rean.	6.8	

The lack of statistically significant future changes is confirmed when using other criteria for the identification of MSWs.



# Type of events: split/displacement MSWs


**Criterion** based on the calculation of area-weighted rotation around the occurrence of MSWs (algorithm developed by K. Shibata).

Model	Past (S/D ratio)	Future (S/D ratio)
CCSRNIES- MIROC3.2	0.13	0.33
NIWA-UKCA	0.50	0.00
GEOS-CCM	0.17	0.40
CNRM-CCM	0.20	0.11
EMAC-O	0.23	0.16
ACCESS	0.21	0.27
NCEP/NCAR rean.	0.77	

- CCMI models show in general a very low number of vortex split MSWs.
- **Future:** no significant changes are found.

# Future seasonal changes in the intensity of PNJ

Daily  $\bar{u}$ @10 hPa (m/s)  
Future-minus-past



- No stat. signif. changes in GEOS-CCM and CNRM-CCM
- Changes in other 4 models:
  - Early winter: stat. signif stronger future PNJ.
  - Midwinter: weaker future PNJ (only stat. signif. in 2 of them).

Contours enclose areas with statistically significant values at a 95% conf. level

# Conclusions

- **No** statistically significant future changes in the mean frequency of MSWs are found in general in the analyzed CCMI models.
- Other characteristics of MSWs do not show stat. significant changes in the future either.
- Two groups of models are found in terms of future seasonal changes in the intensity of the PNJ:
  1. Models that do not show any statistically significant changes of the polar vortex in the future (GEOS-CCM & CNRM-CCM).
  2. Models with a future weakening (strengthening) in midwinter (early winter) (CCSRIES-MIROC3.2, NIWA-UKCA, ACCESS & EMAC-O).



**Thank you for your attention!**